


Quantum Metrology in Superconducting Transmons

Gehad Eldibany

Nordita Summer Internship

Supervisor: Dr Sofia Qvarfort

Motivation: Superconductivity



Schematic of a Josephson junction. Source: Maruf et al. (2018)

- Superconducting quantum circuits with Josephson junctions provide low-dissipation and long-coherence platforms for quantum technologies.
- Transmons are superconducting qubits based on the Josephson cosine potential $V(\hat{\phi}) = -E_J \cos \hat{\phi}$, which introduces nonlinearity into their dynamics and create distinct quantum states (e.g., $|0\rangle$ and $|1\rangle$) for reliable computation.

Motivation: Metrology

- By expanding the cosine potential $V(\hat{\phi})$, we can obtain a Hamiltonian that contains non-linear higher-order terms.
- Nonlinear terms can increase sensitivity to parameters, but the resulting states are more fragile to decoherence, so the practical advantage can vanish if noise dominates.
- An optimal measurement is needed to saturate F_Q (so $F_C = F_Q$) and extract the maximum information available from the noisy quantum state and retain the advantage of non-linear dynamics.

Objective

Achieve $F_C = F_Q$ to maximize metrology precision in transmons.

Quantum and Classical Fisher Information

Quantum Fisher Information (F_Q)

The maximum amount of information about a parameter g obtainable from a quantum state ψ_g through an optimal measurement. Defined as:

$$F_Q = 4 \left(\langle \partial_g \psi_g | \partial_g \psi_g \rangle - |\langle \psi_g | \partial_g \psi_g \rangle|^2 \right) \quad (1)$$

Classical Fisher Information (F_C)

The amount of information about a parameter g obtainable from the probability distribution $p(x|g)$ of a measurement outcome x . Defined as:

$$F_C = \int dx \frac{1}{p(x|g)} \left(\frac{\partial p(x|g)}{\partial g} \right)^2 \quad (2)$$

$F_C \leq F_Q$, equality requires optimal measurement.

Research Questions and Method

To saturate CFI to QFI, we address these questions:

- 1 What is the optimal initial state? \rightarrow *Achieves higher F_Q*
- 2 What is the optimal measurement? \rightarrow *Achieves $F_C = F_Q$*

Method

- 1 Expand the cosine potential to obtain H_{eff} .
- 2 Apply the evolution operator $U(t) = e^{-iH_{\text{eff}}t/\hbar}$ to an initial state.
- 3 Compute F_Q .
- 4 Apply measurement and compute the probability distribution.
- 5 Compute F_C and see how it compares to F_Q for this measurement.

The Effective Hamiltonian

Upon expanding the potential up to the sixth order give:

$$-E_J \cos \hat{\phi} = -E_J + \frac{E_J}{2} \hat{\phi}^2 - \frac{E_J}{24} \hat{\phi}^4 + \frac{E_J}{720} \hat{\phi}^6 - \dots \quad (3)$$

We obtain:

$$H_{\text{eff}} = \hbar\omega_p \hat{n} - \frac{3E_C}{4} \hat{n} - \frac{E_C}{4} \hat{n}(\hat{n} - 1) + \frac{E_C}{18} \sqrt{\frac{2E_C}{E_J}} \hat{n}(\hat{n} - 1)(\hat{n} - 2) \quad (4)$$

Where:

$$\hbar\omega_p = \sqrt{8E_J E_C}, \quad \hat{n} = a^\dagger a, \quad (5)$$

$\hbar\omega_p n$ is the linear energy term, and the remaining terms account for charging energy E_C and Josephson energy E_J (non-linear terms)

Evolution of initial states

- Cat states had a lower performance (i.e. lower F_Q) compared to a superposition of Fock states $|0\rangle + |n\rangle$ when n is large.
- H_{eff} is diagonal in Fock basis n and has eigenvalues

$$E_n = \hbar\omega_p n - \frac{3E_C}{4} n - \frac{E_C}{4} n(n-1) + \frac{E_C}{18} \sqrt{\frac{2E_C}{E_J}} n(n-1)(n-2), \quad (6)$$

with $E_0 = 0$.

- The time-evolved state thus picks a phase:

$$|\psi(t)\rangle = e^{-iH_{\text{eff}}t/\hbar} |\psi_0\rangle = \frac{1}{\sqrt{2}} \left(e^{-iE_0t/\hbar} |0\rangle + e^{-iE_nt/\hbar} |n\rangle \right). \quad (7)$$

$$|\psi(t)\rangle = \frac{1}{\sqrt{2}} \left(|0\rangle + e^{-i\phi} |n\rangle \right), \quad \phi = E_nt/\hbar \quad (8)$$

To calculate F_Q for ω_p , we calculate the partial derivative:

$$\partial_g |\psi\rangle = -itn |\psi\rangle \longrightarrow F_Q(\omega_p) = n^2 t^2 \quad (9)$$

Measurements: Fock VS Homodyne

Fock Measurement

- Counts photon number, $F_C = 0$, not optimal.

Homodyne Measurement


- Measures quadrature (amplitude).
-

$$F_{C_{|0\rangle,|n\rangle}}^{\max} = n^2 t^2 \left[2 - \frac{2}{\sqrt{\pi}} \int_{-\infty}^{\infty} \frac{e^{-x^2}}{1 + \frac{H_n^2(x)}{2^n n!}} dx \right], \quad (10)$$

where $H_n(x)$ are Hermite polynomials.

For $n = 1$ the integral evaluates to 1.15, $n = 2 : 1.30$, $n = 3 : 1.34..$ Thus, $F_C < F_Q$, suboptimal.

Visual Results



Future work

- Try other measurements more optimal for non-linear dynamics, such as the projection into $|+\rangle, |-\rangle$

Acknowledgments

- We thank Dr. Andrea Maiani for inspiring the work on transmon potential.